

## HCU (Ph.D-Physics-2010)

## SECTION - A

Q1. Let

$$f(x) = \begin{cases} \frac{1}{|x|}, & x \geq 1 \\ ax^2 + b, & |x| < 1 \end{cases}$$

For the function  $f(x)$  to be continuous and differentiable at any point,  $a$  and  $b$  should have the values

(a)  $a = 1, b = 0$

(b)  $a = \frac{1}{2}, b = \frac{1}{2}$

(c)  $a = \frac{1}{2}, b = \frac{3}{2}$

(d)  $a = -\frac{1}{2}, b = \frac{3}{2}$

Q2. The sum of the squares of the eigenvalues of  $3 \times 3$  matrix  $A$  is 20. The corresponding sum for the matrix  $2A$  is

(a) 40

(b) 60

(c) 120

(d) 80

Q3. The differential equation and its general solution for the roots of the characteristics equations  $\lambda_1 = 0, \lambda_2 = \lambda_3 = 4$  are given by

(a)  $y''' - 8y'' + 16y' = 0; y = c_1 + (c_2 + c_3x) \cosh 4x$

(b)  $y''' - 8y'' + 16y' = 0; y = c_1 + (c_2 + c_3x) e^{4x}$

(c)  $y''' - 8y'' + 16y' = 0; y = c_1 + (c_2 + c_3x) \sin 4x$

(d)  $y''' - 8y'' + 16y' = 0; y = c_1 + (c_2 + c_3x) e^{4x}$

Q4. If  $\theta$  is real, the function  $\sin(\theta^2)$  is

(a) a periodic function of  $\theta$  with period  $2\pi$ (b) a periodic function of  $\theta$  with period  $(2\pi)^2$ (c) a periodic function of  $\theta$  with period  $\pi^2$ (d) not a periodic function of  $\theta$ **Head office**

fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498

**Branch office**

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16

Q5. One mole of a monatomic ideal gas initially at temperature  $T_0$  expands from volume  $V_0$  to  $2V_0$  at constant pressure. The work of expansion ( $W$ ) and the heat absorbed by the gas ( $Q$ ) are:

(a)  $W = RT_0, Q = \frac{5}{2}RT_0$

(b)  $W = RT_0 \ln 2, Q = RT_0$

(c)  $W = 3RT_0, Q = \frac{3}{2}RT_0$

(d)  $W = \frac{3}{2}RT_0, Q = \frac{5}{2}RT_0$

where  $R$  is the gas constant.

Q6. A body of constant heat capacity  $C_p$  and a temperature  $T_i$  is brought into contact with a reservoir at temperature  $T_f$ . If the equilibrium between the body and the reservoir is established at constant pressure and  $T_i \neq T_f$ , the total entropy change of the body the body plus the reservoir is

(a)  $C_p \ln\left(\frac{T_f}{T_i}\right)$

(b)  $C_p \frac{(T_i - T_f)}{T_f}$

(c)  $C_p \left[ \ln\left(\frac{T_f}{T_i}\right) - 1 + \left(\frac{T_i}{T_f}\right) \right]$

(d)  $C_p \left[ \left(\frac{T_i}{T_f}\right) - \ln\left(\frac{T_f}{T_i}\right) \right]$

Q7. If  $z$  is the canonical partition function of a system, the entropy of the system is

(a)  $k_B \ln Z$

(b)  $k_B T \left( \partial \ln \frac{Z}{\partial V} \right)_T$

(c)  $k_B T \ln z$

(d)  $k_B \left[ \ln Z + T \left( \partial \ln \frac{Z}{\partial T} \right)_P \right]$

Q8. If the interatomic potential energy in a solid is given by  $U(r) = \frac{b}{r^n} - \frac{a}{r^m}$  where  $a$  and  $b$  are positive constants and  $n > m$ , the equilibrium interatomic spacing  $r_0$ , is

(a)  $r_0 = \left[ \left(\frac{a}{b}\right) \left(\frac{m}{n}\right) \right]^{1/(m+n)}$

(b)  $r_0 = \left[ \left(\frac{b}{a}\right) \left(\frac{n}{m}\right) \right]^{1/(n-m)}$

(c)  $r_0 = \left[ \left(\frac{a}{b}\right) \left(\frac{n}{m}\right) \right]^{1/(n-m)}$

(d)  $r_0 = \left[ \left(\frac{b}{a}\right) \left(\frac{m}{n}\right) \right]^{1/(m+n)}$

#### Head office

fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498

#### Branch office

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16

Q9. For non-magnetic metals, specific heat ( $C$ ) at temperatures ( $T \ll \theta_D$ , the Debye temperature) varies with temperature ( $T$ ) as

(a)  $\frac{C}{T} = aT$

(b)  $\frac{C}{T} = a + bT$

(c)  $\frac{C}{T} = a$

(d)  $\frac{C}{T} = a + bT^2$

Q10. A particle of mass  $m$  whose total energy is twice its rest energy collides with an identical particle at rest. If they stick together, the mass  $M$  and the velocity  $v$  of the resulting composite particle are given by

(a)  $M = \sqrt{6}m, v = \frac{c}{\sqrt{3}}$

(b)  $M = \sqrt{3}m, v = c\sqrt{\frac{2}{3}}$

(c)  $M = 2m, v = c\sqrt{\frac{5}{3}}$

(d)  $M = \sqrt{2}m, v = c\sqrt{\frac{7}{3}}$

Q11. In a Compton scattering experiment a photon of energy  $E_0$  is scattered at  $0^\circ, 90^\circ$  and  $180^\circ$  with respect to the incident photon are in the proportion

(a)  $1 : \frac{mc^2}{2E_0 + mc^2} : \frac{mc^2}{E_0 + mc^2}$

(b)  $1 : \frac{mc^2}{E_0 + mc^2} : \frac{mc^2}{2E_0 + mc^2}$

(c)  $1 : \frac{mc^2}{2E_0 + mc^2} : \frac{mc^2}{E_0 - mc^2}$

(d)  $1 : \frac{mc^2}{E_0 + mc^2} : \frac{\sqrt{2}mc^2}{E_0(\sqrt{2}-1) + \sqrt{2}mc^2}$

Q12. Consider 4 infinite sheets of surface charge density as shown in the figure.


The field in region 2 is given by

(a) 0

(b)  $\frac{\sigma}{\epsilon_0} \hat{z}$

(c)  $\frac{-\sigma}{(2\epsilon_0)} \hat{z}$

(d)  $\frac{-\sigma}{\epsilon_0} \hat{z}$




**Head office**

fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498

**Branch office**

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16

Q13. Consider the following charge distributions. All the polygons are regular with edge length  $d$ .



The dipole moment is independent of the choice of origin for configurations


- (a) (ii) and (iii) (b) (i) and (iv)  
(c) (iii) and (iv) (d) (i) and (ii)
- Q14. A hollow metallic waveguide can support  
(a)  $TE, TM$  and  $TEM$  waves (b) Either  $TE$  or  $TM$  waves  
(c) Both  $TE$  and  $TM$  waves (d) only  $TEM$  waves
- Q15. To use a transistor as an amplifier  
(a) emitter-base junction should be forward-biased and the collector base junction reverse biased  
(b) both junctions should be forward biased  
(c) both junctions should be reversed-biased  
(d) it does not matter how the transistor is biased it always works as an amplifier
- Q16. Which of the following logic gates can be used as a controlled inverter?  
(a) AND gate (b) OR gate  
(c) INVERTER gate (d) XOR gate
- Q17. The neutrinos emitted in a  $\beta$  decay process are  
(a) right handed (b) left handed  
(c) both left and right handed (d) spin-zero and massless particles
- Q18. In the fission of  $^{235}\text{U}_92$  the mass ratio of the fission fragment is 1.5. What is the ratio of the velocities of these fragments?  
(a) 3:2 (b) 2:3 (c) 9:4 (d) 4:9

**Head office**

fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498

**Branch office**

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16

- Q19. Alpha decay is an example of
- (a) nuclear fission (b) Barrier penetration  
(c) Coulomb repulsion (d) nuclear ionization
- Q20. Let  $|n\rangle$  denote the normalized eigenstates of the Hamiltonian  $H$  describing a linear harmonic oscillator corresponding to the eigenvalues  $\left(n + \frac{1}{2}\right)\hbar\omega$ . The normalized state  $|\psi\rangle = \frac{1}{\sqrt{5}}[2|1\rangle - |2\rangle]$  is
- (a) an eigenstate of  $H$  corresponding to the eigenvalues  $\frac{1}{2}\hbar\omega$   
(b) an eigenstate of  $H$  corresponding to the eigenvalues  $\frac{11}{2}\hbar\omega$   
(c) an eigenstate of  $H$  corresponding to the eigenvalues 0  
(d) not an eigenstate of  $H$
- Q21.  $\hat{A}, \hat{B}$  and  $\hat{C}$  are three Hermitian operators. Which of the following statements cannot be correct?
- (a)  $[\hat{A}, \hat{B}] = \hat{C}$  (b)  $\{A, B\} = \hat{C}$  (c)  $[A, B] = i\hat{C}$  (d)  $\hat{A}^2 + \hat{B}^2 = \hat{C}^2$
- Q22. The allowed energies for a particle in a one-dimensional square well potential between  $-\alpha$  and  $\alpha$  are
- $$E_n = \frac{\pi^2 \hbar^2 n^2}{8ma^2}$$
- The wave function shown in the figure corresponds to
- 
- (a)  $n=3$  (b)  $n=2$  (c)  $n=4$  (d)  $n=7$
- Q23. With the positive  $z$ -axis chosen to be upwards, the Lagrangian of a particle of mass  $m$  falling under gravity is
- (a)  $\frac{1}{2}m\dot{z}^2 - mgz$  (b)  $-\frac{1}{2}m\dot{z}^2 - mgz$   
(c)  $\frac{1}{2}m\dot{z}^2 + mgz$  (d)  $-\frac{1}{2}m\dot{z}^2 + mgz$

**Head office**


fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498

**Branch office**

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16

- Q24. An aircraft executes a horizontal loop of radius  $1 \text{ km}$  with a steady speed of  $900 \text{ km/h}$ . The ratio of horizontal acceleration of the aircraft to the acceleration due to gravity ( $g = 9.8 \text{ m/sec}^2$ ) is about
- (a) 0                      (b) 5                      (c) 6                      (d) 7
- Q25. Which of the following has non-vanishing Poisson bracket with  $L_z$ , the  $Z$ -component of the angular momentum  $\vec{L}$ ?
- (a)  $a(x^2 + y^2 + z^2)$                       (b)  $a(x^2 + y^2) + bz^2$   $a \neq b$   
 (c)  $ax^2 + by^2 + az^2$   $a \neq b$                       (d)  $a(x^2 + y^2)^2$

## SECTION - B

- Q26. The value of the integral  $\int_{-1}^2 dx x^2 \delta(2x-1)$  where  $\delta(x)$  denotes the Dirac delta function, is
- (a) 0                      (b)  $\frac{1}{4}$                       (c) 1                      (d)  $\frac{1}{8}$
- Q27. If  $I_1$  and  $I_2$  denotes the values of the integral of  $\frac{1}{\left(z + \frac{1}{2}\right)^2}$  evaluated along the semicircle  $C_1$  and  $C_2$  as shown, then  $I_1 - I_2$  is
- (a) 0                      (b)  $2\pi i \times \frac{1}{4}$   
 (c)  $2\pi i \times \frac{1}{8}$                       (d)  $2\pi$
- 
- Q28. For a matrix  $A$  satisfying  $A^3 = 1$ , which of the following cannot be possible set of eigenvalues?
- (a)  $(1, 1, 1)$                       (b)  $\left(1, 1, e^{\frac{2\pi i}{3}}\right)$                       (c)  $\left(1, e^{\frac{2\pi i}{3}}, e^{\frac{4\pi i}{3}}\right)$                       (d)  $\left(1, -e^{\frac{2\pi i}{3}}, e^{\frac{4\pi i}{3}}\right)$

### Head office

fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498

### Branch office

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16

Q29. It is impossible to find a real  $4 \times 4$  matrix with the eigenvalues given by

- (a)  $(1, 1, -1, -1)$       (b)  $(1, i, i, 1)$       (c)  $(1, i - i, 1)$       (d)  $(1, 1, 0, 0)$

Q30. The value of the integral

$$\int_{-\infty}^{\infty} e^{-x^2} H_3(x) P_2(x) dx$$

where  $H_n(x)$  and  $P_n(x)$  are the Hermite and Legendre polynomials, is

- (a)  $2^3 \cdot 3! \sqrt{\pi}$       (b)  $\frac{2}{5}$       (c)  $\sqrt{2^3 \cdot 3! \sqrt{\pi} \cdot \frac{2}{5}}$       (d) 0

Q31. For the function  $f(z) = \frac{z}{(z+2)^2}$  the order of the pole at  $z = -2$  and the residue at the

pole are respectively given by

- (a) 2, -2      (b) 2, 1      (c) 1, -2      (d) 1, 1

Q32. The Bernoulli equation given by  $y' = P(x)y + Q(x)y^m$ ,  $m \neq 0, m \neq 1$ , can be reduced to a linear equation for  $z$  by substitution

- (a)  $z = y^{1-m}$       (b)  $z = y^{m-1}$       (c)  $z = y^{1-2m}$       (d)  $z = y^{2m-1}$

Q33. The rank of the matrix  $\begin{pmatrix} 1 & 1 & -1 & -1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & -1 \end{pmatrix}$  is

- (a) 1      (b) 2      (c) 3      (d) 4

Q34. Let  $|n\rangle$  denotes the eigenstates of linear harmonic oscillator corresponding to the

eigenvalues  $\left(n + \frac{1}{2}\right)\hbar\omega$ . If  $|\psi(0)\rangle = \frac{1}{\sqrt{2}}[|0\rangle + |1\rangle]$  denotes the state of harmonic

oscillator at  $t = 0$  and  $|\psi(t)\rangle$  that at time  $t$ , the  $|\langle\psi(0)|\psi(t)\rangle|^2$  first becomes zero at

- (a)  $t = \frac{\pi}{8\omega}$       (b)  $t = \frac{\pi}{4\omega}$       (c)  $t = \frac{\pi}{2\omega}$       (d)  $t = \frac{\pi}{\omega}$

#### Head office

fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498

#### Branch office

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16



Q35. Which of the following pairs of wave function (with  $A \neq B$ ) describe the same physical state of a free particle

(a)  $\psi_1(x) = Ae^{ikx} + Be^{-ikx}, \psi_2(x) = Ae^{ikx} + Be^{-ikx}$

(b)  $\psi_1(x) = Ae^{ikx} + Be^{-ikx}, \psi_2(x) = Ae^{ikx} - Be^{-ikx}$

(c)  $\psi_1(x) = Ae^{ikx} + Be^{-ikx}, \psi_2(x) = -Ae^{ikx} - Be^{-ikx}$

(d)  $\psi_1(x) = Ae^{ikx} + Be^{-ikx}, \psi_2(x) = Be^{ikx} + Ae^{-ikx}$

Q36. For a spin  $-\frac{1}{2}$  particle in the state  $\begin{pmatrix} \alpha_+ \\ \alpha_- \end{pmatrix}$ , the expectation value of  $\hat{S}_z = \frac{1}{2} \hbar \sigma_z$  is

(a)  $\frac{\hbar}{2} (\alpha_+^* \alpha_- + \alpha_-^* \alpha_+)$

(b)  $\frac{-i\hbar}{2} (\alpha_+^* \alpha_- - \alpha_-^* \alpha_+)$

(c)  $\frac{\hbar}{2} (|\alpha_+|^2 - |\alpha_-|^2)$

(d) 0

Q37. Two spin  $-\frac{1}{2}$  particles with spins  $\hat{S}_1$  and  $\hat{S}_2$  are in a singlet state. The value of the potential  $V(r) = V_1(r) + \frac{1}{\hbar^2} \hat{S}_1 \cdot \hat{S}_2 V_2(r)$ . For such a state will be

(a)  $V_1(r) + \frac{1}{4} V_2(r)$

(b)  $V_1(r) - \frac{3}{4} V_2(r)$

(c) 0

(d)  $V_1(r) - \frac{3}{2} V_2(r)$

Q38. If  $\hat{H}$  and  $\hat{p}$  respectively denotes the Hamiltonian and the momentum operator for a free particle, then the function  $\sin kx$  and  $\cos kx$  are eigenfunctions

(a) of both  $\hat{H}$  and  $\hat{p}$

(b) only of  $\hat{H}$  but not of  $\hat{p}$

(c) only of  $\hat{p}$  but not of  $\hat{H}$

(d) of neither  $\hat{H}$  nor  $\hat{p}$

#### Head office

fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498

#### Branch office

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16



Q39. If the Gibbs free energy of a system is given by  $G(T, P) = RT \ln \left[ \frac{\alpha P}{(RT)^{\frac{5}{2}}} \right]$  where  $\alpha$  and

$R$  are constants, the specific heat at constant pressure,  $C_p$ , is

(a)  $\frac{3}{2}R$

(b)  $\frac{5}{2}R$

(c)  $\frac{5}{2}R - R \ln \left[ \frac{\alpha P}{(RT)^{\frac{5}{2}}} \right]$

(d)  $R \ln \left[ \frac{\alpha P}{(RT)^{\frac{5}{2}}} \right]$

Q40. The entropy of a magnetic system changes by an amount  $\Delta S = \frac{-CH\Delta H}{T^2}$  (where  $C$  is a constant) if its temperature is held constant at  $T$  and the magnetic field is changed from  $H$  to  $H + \Delta H$ . The magnetization  $M$  of such a system depends on temperature  $T$  and field  $H$  as

(a)  $M = \frac{CH^2}{T}$

(b)  $M = \frac{CT}{H}$

(c)  $M = C \left( \frac{T}{H} \right)^2$

(d)  $M = \frac{CH}{T}$

Q41. A system of two energy levels  $\epsilon_0$  and  $\epsilon_1$  ( $\epsilon_1 > \epsilon_0$ ) is populated by  $N$  particle at temperature  $T$  in accordance with the Boltzmann distribution law. The average energy per particle is

(a)  $\epsilon_0 + \epsilon_1 e^{-(\epsilon_1 - \epsilon_0)/k_B T}$

(b)  $\epsilon_1 \pm \epsilon_0 e^{-(\epsilon_1 - \epsilon_0)/k_B T}$

(c)  $\frac{\left[ \epsilon_0 + \epsilon_1 e^{\frac{-(\epsilon_1 - \epsilon_0)}{k_B T}} \right]}{\left[ 1 + e^{\frac{-(\epsilon_1 - \epsilon_0)}{k_B T}} \right]}$

(d)  $\frac{\left[ \epsilon_0 + \epsilon_1 e^{\frac{-(\epsilon_1 - \epsilon_0)}{k_B T}} \right]}{\left[ 1 - e^{\frac{-(\epsilon_1 - \epsilon_0)}{k_B T}} \right]}$

**Head office**

fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498

**Branch office**

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16

Q42. A monatomic gas consists of atoms with two internal energy levels: a ground state of degeneracy  $g_1$  and an excited state of degeneracy  $g_2$  at an energy  $\epsilon$  above the ground state (chosen as the zero of energy scale). The specific heat of the gas is

- (a)  $\frac{3}{2}k_B$  (b)  $\frac{3}{2}k_B - \frac{g_1 e^{-\epsilon/k_B T}}{1 + g_2 e^{-\epsilon/k_B T}}$   
 (c)  $\frac{3}{2}k_B + \frac{g_1 g_2 \epsilon^2 e^{\epsilon/k_B T}}{k_B T^2 (g_2 + g_1 e^{\epsilon/k_B T})^2}$  (d)  $\frac{g_1 g_2 \epsilon^2 e^{-\epsilon/k_B T}}{k_B T (g_2 + g_1 e^{\epsilon/k_B T})^2}$

Q43. Consider an ideal gas consisting of  $N$  particles obeying classical statistics. Suppose that the energy of one particle  $\epsilon$  is proportional to the magnitude of its momentum  $p$  i.e.,  $\epsilon = Cp$ . The partition function of the gas is proportional to

- (a)  $\left(\frac{k_B T}{C}\right)^{3N}$  (b)  $\left(\frac{k_B T}{C}\right)^N$  (c)  $\left(\frac{C}{k_B T}\right)^N$  (d)  $(Ck_B T)^N$

Q44. If in a diffraction experiment,  $X$ -ray photon of wavevector  $\vec{k}$  gets scattered to the final state with wave vector  $\vec{k}' = \vec{k} + \vec{G}$ , where  $\vec{G}$  is a reciprocal lattice vector, the Bragg diffraction condition is given by

- (a)  $\vec{k} \cdot \vec{G} + G^2 = 0$  (b)  $\vec{k} \cdot \vec{G} + G^2 = 0$  (c)  $\vec{k} \cdot \vec{G} = (\vec{G} \cdot \vec{G})/2$  (d)  $\vec{k} \cdot \vec{k} = 2(\vec{k} \cdot \vec{G})$

Q45. If  $\vec{a}_1 = \frac{a}{2}(-\hat{x} + \hat{y} + \hat{z})$ ;  $\vec{a}_2 = \frac{a}{2}(\hat{x} - \hat{y} + \hat{z})$  and  $\vec{a}_3 = \frac{a}{2}(\hat{x} + \hat{y} - \hat{z})$  are the primitive translation vectors of a cubic lattice of size (lattice parameter)  $a$  and  $\hat{x}, \hat{y}, \hat{z}$  are the orthogonal unit vectors parallel to the cube edges, the primitive translation vectors  $\vec{b}_1, \vec{b}_2, \vec{b}_3$  of the reciprocal lattice are given by

- (a)  $\vec{b}_1 = \frac{2\pi}{a}(\hat{y} + \hat{z})$ ;  $\vec{b}_2 = \frac{2\pi}{a}(\hat{z} + \hat{x})$ ;  $\vec{b}_3 = \frac{2\pi}{a}(\hat{x} + \hat{y})$ ;  
 (b)  $\vec{b}_1 = \frac{2\pi}{a}(\hat{y} - \hat{z})$ ;  $\vec{b}_2 = \frac{2\pi}{a}(\hat{z} - \hat{x})$ ;  $\vec{b}_3 = \frac{2\pi}{a}(\hat{x} - \hat{y})$ ;  
 (c)  $\vec{b}_1 = \frac{2\pi}{a}(\hat{y} + \hat{z})$ ;  $\vec{b}_2 = \frac{2\pi}{a}(-\hat{x} + \hat{z})$ ;  $\vec{b}_3 = \frac{2\pi}{a}(-\hat{x} + \hat{y})$ ;  
 (d)  $\vec{b}_1 = \frac{2\pi}{a}(-\hat{y} + \hat{z})$ ;  $\vec{b}_2 = \frac{2\pi}{a}(-\hat{z} + \hat{x})$ ;  $\vec{b}_3 = \frac{2\pi}{a}(-\hat{x} + \hat{y})$ ;

### Head office

fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498

### Branch office

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16

Q46. If the atoms vibrating in a lattice at finite temperature are considered to be a collection of  $N$  identical harmonic oscillators, as Einstein did, the heat capacity at constant volume for such a solid is given by

(a)  $C_V = Nk_B x \frac{e^x}{e^x - 1}$                       (b)  $C_V = Nk_B x^2 \frac{e^x}{e^x - 1}$

(c)  $C_V = Nk_B x^2 \frac{e^x}{(e^x - 1)^2}$                       (d)  $C_V = Nk_B x \frac{e^x}{(e^x - 1)^2}$

Q47. Consider a system of  $N$  weakly interacting particles, each of spin  $\frac{1}{2}$  and magnetic moment  $\mu_B$ , exposed to an external magnetic field  $H$ . If the system is in thermal contact with a heat reservoir at temperature  $T$ , the mean energy of the system as a function of  $T$  and  $H$  is given by


(a)  $N \mu_B \sinh(\mu_B H / k_B T)$                       (b)  $N \mu_B \coth(\mu_B H / k_B T)$

(c)  $N \mu_B \tanh^{-1}(\mu_B H / k_B T)$                       (d)  $N \mu_B \tanh(\mu_B H / k_B T)$

Q48. Which of the following relations is consistent with the London equation  $\text{curl } \vec{j} = -\frac{c}{4\pi\lambda_L^2} \vec{B}$  for a superconductor, where  $\lambda_L$  is the London penetration depth.

(a)  $\nabla \vec{B} = \vec{B} / \lambda_L$                       (b)  $\nabla^2 \vec{B} = \vec{B} / \lambda_L$                       (c)  $\nabla^2 \vec{B} = \vec{B} / \lambda_L^2$                       (d)  $\nabla \vec{B} = \vec{B} / \lambda_L^2$

Q49. The output  $V_0$  of the circuit is



(a)  $V_0 = 0$

(b)  $V_0 = (R_2 / R_1) V_1$

(c)  $V_0 = (1 + R_2 / R_1) V_1$

(d)  $V_0 = -(R_2 / R_1) V_1$

**Head office**

fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498


**Branch office**

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16

Q50. In the following figure, the transistor has  $\beta_{dc}$  of 80 and  $V_{CE(sat)}$  of 0.1V .

$R_B$  is adjusted to get transistor into saturation. The value of  $I_{C(sat)}$  and the corresponding value of  $R_B$  are

- (a) 114 mA and 17 k $\Omega$  (b) 1.43 mA and 17 k $\Omega$   
 (c) 114 mA and 1.7 k $\Omega$  (d) 1.43 mA and 17.48 k $\Omega$



Q51. In emitter follower configuration of a transistor

- (a) the emitter current follows the collector current and both are in same phase  
 (b) the collector current follows the emitter current and both are in same phase  
 (c) the base current follows the emitter current and is in opposite phase  
 (d) the emitter current follows the base current and both are in same phase

Q52. Binary equivalent of  $(8.2)_{10}$  is

- (a)  $(1001.0011)_2$  (b)  $(1000.0011)_2$   
 (c)  $(1000.0110)_2$  (d)  $(0111.0110)_2$

Q53. In a pure germanium the intrinsic concentration of charge ( $n_i$ ) at room temperature is  $2.5 \times 10^{13} \text{ cm}^{-3}$ . The electron and hole mobilities are given by  $\mu_n = 3800 \text{ cm}^2/\text{V-s}$  and  $\mu_p = 1800 \text{ cm}^2/\text{V-s}$  respectively. The resistivity of the intrinsic germanium is

- (a) 22.3 $\Omega\text{-cm}$  (b) 66.9 $\Omega\text{-cm}$  (c) 44.6 $\Omega\text{-cm}$  (d) 11.2 $\Omega\text{-cm}$

Q54. The change in isospin ( $\Delta I$ ) in the process  $\Lambda^0 \rightarrow p + \pi^-$  is

- (a)  $\Delta I = 0, 1$  (b)  $\Delta I = \frac{1}{2}, \frac{3}{2}$  (c)  $\Delta I = 0, \frac{1}{2}$  (d)  $\Delta I = 1, 2$

Q55. A spin-0 particle A decaying in its rest frame into particles B and C, through strong interaction. Which of the following statement is not true?

- (a)  $0^+ \rightarrow 0^+ + 0^+$  (b)  $0^+ \rightarrow 0^- + 0^-$  (c)  $0^- \rightarrow 0^+ + 0^-$  (d)  $0^- \rightarrow 0^- + 0^-$

where  $J^p = 0^+$  or  $0^-$  representing the standard convention for labeling the spin and intrinsic parity of a particle.

**Head office**

fiziks, H.No. 40-D, G.F., Jia Sarai,  
 Near IIT, Hauz Khas, New Delhi-16  
 Phone: 011-26865455/+91-9871145498

**Branch office**

Anand Institute of Mathematics,  
 28-B/6, Jia Sarai, Near IIT  
 Hauz Khas, New Delhi-16

- Q56. Which of the following reactions represents a strong interaction
- (a)  $\pi^- + p \rightarrow \Lambda + K^0$  (b)  $\rho^0 \rightarrow \pi^0 + \pi^0$   
 (c)  $K^- \rightarrow \pi^- + \pi^0$  (d)  $p + \bar{p} \rightarrow \pi^+ + \pi^0$
- Q57. In the nuclear shell model,  ${}^{16}_8\text{O}$  is a good closed-shell nucleus and has spin parity  $J^P = 0^+$ . What are the predicted  $J^P$  values for  ${}^{15}_8\text{O}$  and  ${}^{17}_8\text{O}$ ?
- (a)  $0^+, 0^+$  (b)  $\frac{1^-}{2}, \frac{5^+}{2}$  (c)  $\frac{3^-}{2}, \frac{5^+}{2}$  (d)  $0^+, \frac{1^+}{2}$
- Q58. a pion at rest decays to a muon and a neutrino. The magnitude of momentum of the outgoing muon is
- (a)  $\frac{m_\pi^2 - m_\mu^2}{2m_\pi} \cdot c$  (b)  $\frac{m_\pi^2 + m_\mu^2}{2m_\pi} \cdot c$  (c)  $\frac{m_\pi^2 + m_\mu^2}{2m_\mu} \cdot c$  (d)  $\frac{(m_\pi - m_\mu)^2}{2m_\pi} \cdot c$
- Q59. An experiment is performed to search for the evidence of the particle  $H$  through the strong reaction  $p + p \rightarrow H + K^+ + K^+$ . The electric charge ( $Q$ ), Baryon number ( $B$ ) and strangeness quantum number ( $S$ ) of the particle  $H$  will be
- (a)  $Q = 0, B = 0, S = -2$  (b)  $Q = 0, B = 2, S = -2$   
 (c)  $Q = 0, B = 2, S = 2$  (d)  $Q = 0, B = 2, S = 0$
- Q60. Let a plane monochromatic wave be totally internally reflected at the interface  $z = 0$  between two dielectrics. Let  $x-z$  be the plane of incidence with  $x$  along the interface. The planes of constant amplitudes and constant phases in medium 2 are given by
- (a)  $z = c_1, x = c_2$  (b)  $x + z = c_1, x - z = c_2$   
 (c)  $x = c_1, z = c_2$  (d)  $x - z = c_1, x + z = c_2$
- (here  $c_1$  and  $c_2$  are constant)

**Head office**

fiziks, H.No. 40-D, G.F., Jia Sarai,  
 Near IIT, Hauz Khas, New Delhi-16  
 Phone: 011-26865455/+91-9871145498

**Branch office**

Anand Institute of Mathematics,  
 28-B/6, Jia Sarai, Near IIT  
 Hauz Khas, New Delhi-16

Q61. The transformation that reduces the wave equation

$$\frac{\partial^2 u(x,t)}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2 u(x,t)}{\partial t^2} = 0$$

to the form  $\frac{\partial^2 u(\xi, \tau)}{\partial \xi \partial \tau} = 0$  is given by

- (a)  $\xi = c/(ct+x)$ ,  $\tau = c/(ct-x)$       (b)  $\xi = it+x/c$ ,  $\tau = it-x/c$  where  $i = \sqrt{-1}$   
 (c)  $\xi = t+x/c$ ,  $\tau = t-x/c$       (d)  $\xi = t+ix/c$ ,  $\tau = t-ix/c$  where  $i = \sqrt{-1}$

Q62. In the domain of anomalous dispersion, the group velocity ( $v_g$ ), refractive index ( $n$ ) and speed of light in vacuum ( $c$ ) obey

- (a)  $v_g > c/n$       (b)  $v_g = c/n$       (c)  $v_g < c/n$       (d)  $c/n < v_g < c$

Q63. The resolving power of a grating of a given width for a given order of diffraction can be increased by

- (a) making thinner grooves without changing the groove separation  
 (b) making thinner grooves and proportionately decreasing the groove separation  
 (c) making wider grooves without changing the groove separation  
 (d) making wider grooves and proportionately increasing the groove separation

Q64. In a Newton's ring experiment, a plano-convex lens with radius of curvature of the convex surface  $4.18m$  is placed on an optically flat surface with its convex surface down. Interference fringes are observed using a traveling microscope and sodium light ( $\lambda = 589.3nm$ ). The radius of the tenth dark ring is

- (a)  $4.96mm$       (b)  $6.92mm$       (c)  $1.63mm$       (d)  $2.45mm$

Q65. Consider a uniformly charged sphere of radius  $R$  with total charge  $q$ . Let the sphere be coated with a medium with charge density  $\rho = \alpha/r$  (where  $\alpha > 0, r \geq R$ ). The electric field  $\vec{E}$  outside the charged sphere will be independent of  $r$  when

- (a)  $q = 2\pi\alpha R^2 / \epsilon_0$ ,  $\vec{E} = \alpha / (2\epsilon_0) \hat{r}$       (b)  $q = 2\pi\alpha R^2$ ,  $\vec{E} = \alpha / (\epsilon_0) \hat{r}$   
 (c)  $q = \pi\alpha R^2$ ,  $\vec{E} = \alpha / (2\epsilon_0) \hat{r}$       (d)  $q = 2\pi\alpha R^2$ ,  $\vec{E} = \alpha / (2\epsilon_0) \hat{r}$




#### Head office

fiziks, H.No. 40-D, G.F., Jia Sarai,  
 Near IIT, Hauz Khas, New Delhi-16  
 Phone: 011-26865455/+91-9871145498

#### Branch office

Anand Institute of Mathematics,  
 28-B/6, Jia Sarai, Near IIT  
 Hauz Khas, New Delhi-16


Q66. Consider two infinite current carrying wires rigidly held parallel to the  $z$ -axis. The current through the wires is  $I$ . The distance between the wires is  $d$ . The magnetic field at the points marked  $P$  in the three cases shown in the figure below are



- (a) (i)  $\frac{2\mu_0 I}{\pi d} \hat{y}$ ; (ii) 0; (iii)  $-\frac{4\mu_0 I}{3\pi d} \hat{y}$       (b) (i) 0; (ii)  $-\frac{2\mu_0 I}{\pi d} \hat{y}$ ; (iii)  $\frac{4\mu_0 I}{3\pi d} \hat{y}$   
 (c) (i) 0; (ii)  $\frac{2\mu_0 I}{\pi d} \hat{y}$ ; (iii)  $-\frac{4\mu_0 I}{3\pi d} \hat{y}$       (d) (i)  $-\frac{2\mu_0 I}{\pi d} \hat{y}$ ; (ii) 0; (iii)  $-\frac{4\mu_0 I}{3\pi d} \hat{y}$

Q67. Two infinite conducting planes intersect at the origin (see figure). All distances are in meters.

A charge  $q$  is placed at  $(2,1)$ . The potential at the point  $(1,2)$  will be



- (a)  $\frac{q}{4\pi \epsilon_0} \frac{4\sqrt{5}-6}{3\sqrt{10}}$       (b)  $\frac{q}{4\pi \epsilon_0} \frac{2\sqrt{5}-6}{3\sqrt{10}}$   
 (c)  $\frac{q}{4\pi \epsilon_0} \frac{4\sqrt{5}+6}{3\sqrt{10}}$       (d)  $\frac{q}{4\pi \epsilon_0} \frac{-2\sqrt{5}-6}{3\sqrt{10}}$

Q68. The mast of a sailboat makes an angle  $\theta$  with respect to its deck. An observer standing on the dock sees the boat go by with a speed  $v$ . The angle made by the mast as seen by the observer is

- (a)  $\tan^{-1}\left(\sqrt{1-v^2/c^2} \tan \theta\right)$       (b)  $\tan^{-1}\left(\frac{\tan \theta}{\sqrt{1-v^2/c^2}}\right)$   
 (c)  $\tan^{-1}\left(\tan\left(\theta\sqrt{1-v^2/c^2}\right)\right)$       (d)  $\tan^{-1}\left(\tan\left(\theta/\sqrt{1-v^2/c^2}\right)\right)$

**Head office**

fiziks, H.No. 40-D, G.F., Jia Sarai,  
 Near IIT, Hauz Khas, New Delhi-16  
 Phone: 011-26865455/+91-9871145498

**Branch office**

Anand Institute of Mathematics,  
 28-B/6, Jia Sarai, Near IIT  
 Hauz Khas, New Delhi-16



Q69. The coherence matrix for light propagating along  $z$  - direction can be written down as

$$I_{xy} = \begin{pmatrix} \overline{E_x E_x^*} & \overline{E_x E_y^*} \\ \overline{E_y E_x^*} & \overline{E_y E_y^*} \end{pmatrix}$$

where  $E_x$  and  $E_y$  are the projections of the electric field along the  $x$  and  $y$  axes and the overbars denote averaging over time for long enough observation periods. If the determinant of the matrix equals  $I^2/4$  (where  $I = \overline{|E_x|^2} + \overline{|E_y|^2}$ ) then the light is

- (a) completely polarized (b) completely unpolarized  
(c) 50% partially polarized (d) 25% partially polarized

Q70. For a particle in a one dimensional potential

$$V(x) = -V_0 \operatorname{sech}^2(\alpha x)$$

the time period of small oscillations about the equilibrium is

- (a)  $2\pi\sqrt{\frac{m}{2V_0\alpha^2}}$  (b)  $2\pi\sqrt{\frac{V_0}{m\alpha^2}}$  (c)  $2\pi\sqrt{\frac{1}{m\alpha^2V_0}}$  (d)  $2\pi\sqrt{\frac{m}{V_0\alpha^2}}$

Q71. The Lagrangian of a point particle is

$$L = \frac{1}{2}m\dot{x}^2 - \frac{1}{2}\omega^2x^2 - \alpha x^4 + \beta x\dot{x}^2$$

Its equation of motion is

- (a)  $m\ddot{x} + \beta\dot{x}^2 + \omega^2x + 4\alpha x^3 = 0$  (b)  $m\ddot{x} - \beta\dot{x}^2 - \omega^2x + 4\alpha x^3 = 0$   
(c)  $(m + 2\beta x)\ddot{x} + \beta\dot{x}^2 + \omega^2x + 4\alpha x^3 = 0$  (d)  $(m + \beta x)\ddot{x} + \beta\dot{x}^2 + \omega^2x + 4\alpha x^3 = 0$

Q72. A particle of mass  $m$  moves in a spherically symmetric potential  $V(r) = \frac{1}{2}kr^2$  and has angular momentum  $L$ . Which of the following is a constant of motion?

- (a)  $\frac{1}{2}m\left(\frac{d\vec{r}}{dt}\right)^2 + \frac{1}{2}kr^2 + \frac{L^2}{2mr^2}$  (b)  $\frac{1}{2}m\left(\frac{dr}{dt}\right)^2 + \frac{1}{2}kr^2$   
(c)  $\frac{1}{2}m\left(\frac{d\vec{r}}{dt}\right)^2 + \frac{L^2}{2mr^2}$  (d)  $\frac{1}{2}m\left(\frac{dr}{dt}\right)^2 + \frac{1}{2}kr^2 + \frac{1}{2}\frac{L^2}{2mr^2}$

#### Head office

fiziks, H.No. 40-D, G.F., Jia Sarai,  
Near IIT, Hauz Khas, New Delhi-16  
Phone: 011-26865455/+91-9871145498

#### Branch office

Anand Institute of Mathematics,  
28-B/6, Jia Sarai, Near IIT  
Hauz Khas, New Delhi-16

Q73. A particle of mass  $m$  and angular momentum  $L^2 = 10mV_0R^2$  moves in a potential

$$V(\vec{r}) = -V_0 \left( \frac{3R}{r} + \frac{R^3}{r^3} \right), V_0 > 0. \text{ The radius of the stable circular orbit is}$$

- (a)  $R$                       (b)  $R/2$                       (c)  $2R$                       (d)  $3R$


Q74. The Poisson bracket of  $L_x$  and  $\vec{p}^2$ , where  $\vec{L} = \vec{r} \times \vec{p}$  denotes the angular momentum, is

- (a)  $2L_x$                       (b)  $2p_x$                       (c)  $L_x p_x$                       (d) 0

Q75. Consider a diatomic molecule undergoing a vibrational electronic transition from  $V''$  to  $V'$ .

Which one of the following is correct?

- (a) The most intense line will be (0,0) transition  
 (b) The most intense line will be (0,1) transition  
 (c) The most intense line will be (0,2) transition  
 (d) The most intense line will be (0,3) transition



#### Head office

fiziks, H.No. 40-D, G.F., Jia Sarai,  
 Near IIT, Hauz Khas, New Delhi-16  
 Phone: 011-26865455/+91-9871145498

#### Branch office

Anand Institute of Mathematics,  
 28-B/6, Jia Sarai, Near IIT  
 Hauz Khas, New Delhi-16