


JEST 2017

Part-A: 1-Mark Questions

Q1. A thin air film of thickness d is formed in a glass medium. For normal incidence, the condition for constructive interference in the reflected beam is (in terms of wavelength λ and integer $m = 0, 1, 2, \dots$)

- (a) $2d = (m - 1/2)\lambda$ (b) $2d = m\lambda$
 (c) $2d = (m - 1)\lambda$ (d) $2\lambda = (m - 1/2)d$

Q2. Consider the circuit shown in the figure where $R_1 = 2.07 k\Omega$ and $R_2 = 1.93 k\Omega$. Current source I delivers $10 mA$ current. The potential across the diode D is $0.7 V$. What is the potential at A ?



- (a) $10.35 V$ (b) $9.65 V$ (c) $19.30 V$ (d) $4.83 V$

Q3. $\int_{-\infty}^{+\infty} (x^2 + 1) \delta(x^2 - 3x + 2) dx = ?$

- (a) 1 (b) 2 (c) 5 (d) 7

Q4. A bead of mass M slides along a parabolic wire described by $z = 2(x^2 + y^2)$. The wire rotates with angular velocity Ω about the z -axis. At what value of Ω does the bead maintain a constant nonzero height under the action of gravity along $-\hat{z}$?

- (a) $\sqrt{3g}$ (b) \sqrt{g} (c) $\sqrt{2g}$ (d) $\sqrt{4g}$

Q5. Which one is the image of the complex domain $\{z | xy \geq 1, x + y > 0\}$ under the mapping

$f(z) = z^2$, if $z = x + iy$?

- (a) $\{z | xy \geq 1, x + y > 0\}$ (b) $\{z | x \geq 2, x + y > 0\}$
 (c) $\{z | y \geq 2\forall x\}$ (d) $\{z | y \geq 1\forall x\}$

Head office


fiziks, H.No. 40 D, G.F, Jia Sarai,
 Near IIT, Hauz Khas, New Delhi-16
 Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
 28-B/6, Jia Sarai, Near IIT
 Hauz Khas, New Delhi-16

- Q6. After the detonation of an atom bomb, the spherical ball of gas was found to be of 15 meter radius at a temperature of $3 \times 10^5 K$. Given the adiabatic expansion coefficient $\gamma = 5/3$, what will be the radius of the ball when its temperature reduces to $3 \times 10^3 K$?
- (a) $156m$ (b) $50m$ (c) $150m$ (d) $100m$

- Q7. What is Y for the circuit shown below?



- (a) $Y = \overline{(A + \bar{B})}(\bar{B} + C)$ (b) $Y = \overline{(A + \bar{B})}(B + C)$
- (c) $Y = \overline{(\bar{A} + B)}(\bar{B} + C)$ (d) $Y = \overline{(A + B)}(\bar{B} + C)$
- Q8. What is the dimension of $\frac{\hbar \partial \psi}{i \partial x}$, where ψ is a wavefunction in two dimensions?
- (a) $kg m^{-1} s^{-2}$ (b) $kg s^{-2}$ (c) $kg m^2 s^{-2}$ (d) $kg s^{-1}$
- Q9. A plane electromagnetic wave propagating in air with $\vec{E} = (8\hat{i} + 6\hat{j} + 5\hat{k})e^{i(\omega t + 3x - 4y)}$ is incident on a perfectly conducting slab positioned at $x = 0$. \vec{E} field of the reflected wave is
- (a) $(-8\hat{i} - 6\hat{j} - 5\hat{k})e^{i(\omega t + 3x + 4y)}$ (b) $(-8\hat{i} + 6\hat{j} - 5\hat{k})e^{i(\omega t + 3x + 4y)}$
- (c) $(-8\hat{i} + 6\hat{j} - 5\hat{k})e^{i(\omega t - 3x - 4y)}$ (d) $(-8\hat{i} - 6\hat{j} - 5\hat{k})e^{i(\omega t - 3x - 4y)}$
- Q10. Let $\Lambda = \begin{pmatrix} 1 & 0 \\ 0 & 11 \end{pmatrix}$ and $M = \begin{pmatrix} 10 & 3i \\ -3i & 2 \end{pmatrix}$. Similarly, transformation of M to Λ can be performed by
- (a) $\frac{1}{\sqrt{10}} \begin{pmatrix} 1 & 3i \\ 3i & 1 \end{pmatrix}$ (b) $\frac{1}{\sqrt{9}} \begin{pmatrix} 1 & -3i \\ 3i & 11 \end{pmatrix}$

Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

(c) $\frac{1}{\sqrt{10}} \begin{pmatrix} 1 & 3i \\ -3i & 11 \end{pmatrix}$

(d) $\frac{1}{\sqrt{9}} \begin{pmatrix} 1 & 3i \\ -3i & 1 \end{pmatrix}$

Q11. (Q_1, Q_2, P_1, P_2) and (q_1, q_2, p_1, p_2) are two sets of canonical coordinates, where Q_i and q_i are the coordinates and P_i and p_i are the corresponding conjugate momenta. If $P_1 = q_2$ and $P_2 = p_1$, then which of the following relations is true?

(a) $Q_1 = q_1, Q_2 = p_2$

(b) $Q_1 = p_2, Q_2 = q_1$

(c) $Q_1 = -p_2, Q_2 = q_1$

(d) $Q_1 = q_1, Q_2 = -p_2$

Q12. Consider magnetic vector potential \vec{A} and scalar potential Φ which define the magnetic field \vec{B} and electric field \vec{E} . If one adds $-\vec{\nabla}\lambda$ to \vec{A} for a well-defined λ , then what should be added to Φ so that \vec{E} remains unchanged up to an arbitrary function of time, $f(t)$?


(a) $\frac{\partial \lambda}{\partial t}$

(b) $-\frac{\partial \lambda}{\partial t}$

(c) $\frac{1}{2} \frac{\partial \lambda}{\partial t}$

(d) $-\frac{1}{2} \frac{\partial \lambda}{\partial t}$

Q13. In the following silicon diode circuit ($V_B = 0.7V$), determine the output voltage waveform (V_{out}) for the given input wave.




Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16




Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16



Q14. $\phi_0(x)$ and $\phi_1(x)$ are respectively are orthonormal wavefunctions of the ground and first excited states of a one dimensional simple harmonic oscillator. Consider the normalised wave function $\psi(x) = c_0\phi_0(x) + c_1\phi_1(x)$, where c_0 and c_1 are real. For what values of c_0 and c_1 will $\langle \psi(x)|x|\psi(x) \rangle$ be maximized?

(a) $c_0 = c_1 = +1/\sqrt{2}$

(b) $c_0 = -c_1 = +1/\sqrt{2}$

(c) $c_0 = +\sqrt{3}/2, c_1 = +1/2$

(d) $c_0 = +\sqrt{3}/2, c_1 = -1/2$


Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

- Q15. Consider the following circuit in steady state condition. Calculate the amount of charge stored in $1\mu F$ and $2\mu F$ capacitors respectively.



- (a) $4\mu C$ and $8\mu C$ (b) $8\mu C$ and $4\mu C$
 (c) $3\mu C$ and $6\mu C$ (d) $6\mu C$ and $3\mu C$
- Q16. If the mean square fluctuations in energy of a system in equilibrium at temperature T is proportional to T^α , then the energy of the system is proportional to
- (a) $T^{\alpha-2}$ (b) $T^{\frac{\alpha}{2}}$ (c) $T^{\alpha-1}$ (d) T^α
- Q17. Suppose the spin degrees of freedom of a 2-particle system can be described by a 21-dimensional Hilbert subspace. Which among the following could be the spin of one of the particles?
- (a) $\frac{1}{2}$ (b) 3 (c) $\frac{3}{2}$ (d) 2
- Q18. Water is poured at a rate of $Rm^3/hour$ from the top into a cylindrical vessel of diameter D . The vessel has a small opening of area a ($\sqrt{a} \ll D$) at the bottom. What should be the minimum height of the vessel so that water does not overflow?
- (a) ∞ (b) $\frac{R^2}{2ga^2}$ (c) $\frac{R^2}{2gaD^2}$ (d) $\frac{8R^2}{\pi D^2 g^2}$

Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
 Near IIT, Hauz Khas, New Delhi-16
 Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
 28-B/6, Jia Sarai, Near IIT
 Hauz Khas, New Delhi-16

Q19. Suppose that we toss two fair coins hundred times each. The probability that the same number of heads occur for both coins at the end of the experiment is

(a) $\left(\frac{1}{4}\right)^{100} \sum_{n=0}^{100} \binom{100}{n}$

(b) $2\left(\frac{1}{4}\right)^{100} \sum_{n=0}^{100} \binom{100}{n}^2$

(c) $\frac{1}{2}\left(\frac{1}{4}\right)^{100} \sum_{n=0}^{100} \binom{100}{n}^2$

(d) $\left(\frac{1}{4}\right)^{100} \sum_{n=0}^{100} \binom{100}{n}^2$

Q20. What is the equation of the plane which is tangent to the surface $xyz = 4$ at the point $(1, 2, 2)$?

(a) $x + 2y + 4z = 12$

(b) $4x + 2y + z = 12$

(c) $x + 4y + z = 0$

(d) $2x + y + z = 6$

Q21. If the ground state wavefunction of a particle moving in a one dimensional potential is proportional to $\exp(-x^2/2)\cosh(\sqrt{2}x)$, then the potential in suitable units such that $\hbar = 1$, is proportional to

(a) x^2

(b) $x^2 - 2\sqrt{2}x \tanh(\sqrt{2}x)$

(c) $x^2 - 2\sqrt{2}x \tan(\sqrt{2}x)$

(d) $x^2 - 2\sqrt{2}x \coth(\sqrt{2}x)$

Q22. A possible Lagrangian for a free particle is

(a) $L = \dot{q}^2 - q^2$ (b) $L = \dot{q}^2 - q\dot{q}$ (c) $L = \dot{q}^2 - q$ (d) $L = \dot{q}^2 - \frac{1}{q}$

Q23. A rod of mass m and length l is suspended from two massless vertical springs with a spring constants k_1 and k_2 . What is the Lagrangian for the system, if x_1 and x_2 be the displacements from equilibrium position of the two ends of the rod?

(a) $\frac{m}{8}(\dot{x}_1^2 + 2\dot{x}_1\dot{x}_2 + \dot{x}_2^2) - \frac{1}{2}k_1x_1^2 - \frac{1}{2}k_2x_2^2$

(b) $\frac{m}{2}(\dot{x}_1^2 + \dot{x}_1\dot{x}_2 + \dot{x}_2^2) - \frac{1}{4}(k_1 + k_2)(x_1^2 + x_2^2)$

(c) $\frac{m}{6}(\dot{x}_1^2 + \dot{x}_1\dot{x}_2 + \dot{x}_2^2) - \frac{1}{2}k_1x_1^2 - \frac{1}{2}k_2x_2^2$

(d) $\frac{m}{2}(\dot{x}_1^2 - 2\dot{x}_1\dot{x}_2 + \dot{x}_2^2) - \frac{1}{4}(k_1 - k_2)(x_1^2 + x_2^2)$

Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

Q24. Two equal positive charges of magnitude $+q$ separated by a distance d are surrounded by a uniformly charged thin spherical shell of radius $2d$ bearing a total charge $-2q$ and centred at the midpoint between the two positive charges. The net electric field at distance τ from the midpoint ($\gg d$) is

- (a) zero (b) proportional to d
 (c) proportional to $1/r^3$ (d) proportional to $1/r^4$

Q25. If the Hamiltonian of a classical particles is $H = \frac{p_x^2 + p_y^2}{2m} + xy$, then $\langle x^2 + xy + y^2 \rangle$ at temperature T is equal to

- (a) $k_B T$ (b) $\frac{1}{2} k_B T$ (c) $2k_B T$ (d) $\frac{3}{2} k_B T$

Part-B: 3-Mark Questions

Q1. A solid, insulating sphere of radius 1cm has charge 10^{-7}C distributed uniformly over its volume. It is surrounded concentrically by a conducting thick spherical shell of inner radius 2cm , outer radius 2.5cm and is charged with $-2 \times 10^{-7}\text{C}$. What is the electrostatic potential in Volts on the surface of the sphere?

Q2. A particle is described by the following Hamiltonian

$$\hat{H} = \frac{\hat{p}^2}{2m} + \frac{1}{2} m \omega^2 \hat{x}^2 + \lambda \hat{x}^4$$

where the quartic term can be treated perturbatively. If ΔE_0 and ΔE_1 denote the energy correction of $O(\lambda)$ to the ground state and the first excited state respectively, what is the fraction $\Delta E_1 / \Delta E_0$?


Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
 Near IIT, Hauz Khas, New Delhi-16
 Phone: 011-26865455/+91-9871145498


Branch office

Anand Institute of Mathematics,
 28-B/6, Jia Sarai, Near IIT
 Hauz Khas, New Delhi-16

- Q3. A simple pendulum has a bob of mass 1 kg and charge 1 Coulomb . It is suspended by a massless string of length 13 m . The time period of small oscillations of this pendulum is T_0 . If an electric field $\vec{E} = 100\hat{x}\text{ V/m}$ is applied, the time period becomes T . What is the value of $(T_0/T)^4$?



- Q4. Let a particle of mass $1 \times 10^{-9}\text{ kg}$, constrained to have one dimensional motion, be initially at the origin ($x = 0\text{ m}$). The particle is in equilibrium with a thermal bath ($k_B T = 10^{-8}\text{ J}$). What is $\langle x^2 \rangle$ of the particle after a time $t = 5\text{ s}$?
- Q5. For the circuit shown below, what is the ratio $\frac{I_1}{I_2}$?



- Q6. A ball of mass 0.1 kg and density 2000 kg/m^3 is suspended by a massless string of length 0.5 m under water having density 1000 kg/m^3 . The ball experiences a drag force, $\vec{F}_d = -0.2(\vec{v}_b - \vec{v}_w)$, where \vec{v}_b and \vec{v}_w are the velocities of the ball and water respectively. What will be the frequency of small oscillations for the motion of pendulum, if the water is at rest?

Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

- Q7. Suppose that the number of microstates available to a system of N particles depends on N and the combined variable UV^2 , where U is the internal energy and V is the volume of the system. The system initially has volume $2m^3$ and energy $200J$. It undergoes an isentropic expansion to volume $4m^3$. What is the final pressure of the system in SI units?
- Q8. The temperature in a rectangular plate bounded by the lines, $x=0, y=0, x=3$ and $y=5$ is $T = xy^2 - x^2y + 100$. What is the maximum temperature difference between two points on the plate?
- Q9. A sphere of inner radius $1cm$ and outer radius $2cm$, centered at origin has a volume charge density $\rho_0 = \frac{K}{4\pi r}$, where K is a nonzero constant and r is the radial distance. A point charge of magnitude $10^{-3}C$ is placed at the origin. For what value of K in units of C/m^2 the electric field inside shell is constant?
- Q10. If $\hat{x}(t)$ be the position operator at a time t in the Heisenberg picture for a particle described by the Hamiltonian, $\hat{H} = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega^2\hat{x}^2$ what is $e^{i\omega t} \langle 0 | \hat{x}(t) \hat{x}(0) | 0 \rangle$ in units of $\frac{\hbar}{2m\omega}$ where $|0\rangle$ is the ground state?

Part-C: 3-Mark Questions

- Q1. Consider a grounded conducting plane which is infinitely extended perpendicular to the y -axis at $y=0$. If an infinite line of charge per unit length λ runs parallel to x -axis at $y=d$, then surface charge density on the conducting plane is
- (a) $\frac{-\lambda d}{(x^2 + d^2 + z^2)}$ (b) $\frac{-\lambda d}{(x^2 + d^2 + z^2)}$
- (c) $\frac{-\lambda d}{\pi(x^2 + d^2 + z^2)}$ (d) $\frac{-\lambda d}{2\pi(x^2 + d^2 + z^2)}$

Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

Q2. A system of particles on N lattice sites is in equilibrium at temperature T and chemical potential μ . Multiple occupancy of the sites is forbidden. The binding energy of a particle at each site is $-\epsilon$. The probability of no site being occupied is

(a) $\frac{1 - e^{\beta(\mu + \epsilon)}}{1 - e^{(N+1)\beta(\mu + \epsilon)}}$

(b) $\frac{1}{[1 + e^{\beta(\mu + \epsilon)}]^N}$

(c) $\frac{1}{[1 + e^{-\beta(\mu + \epsilon)}]^N}$

(d) $\frac{1 - e^{\beta(\mu + \epsilon)}}{1 - e^{-(N+1)\beta(\mu + \epsilon)}}$

Q3. The integral $I = \int_1^\infty \frac{\sqrt{x-1}}{(1+x)^2} dx$ is


(a) $\frac{\pi}{\sqrt{2}}$

(b) $\frac{\pi}{2\sqrt{2}}$

(c) $\frac{\sqrt{\pi}}{2}$

(d) $\sqrt{\frac{\pi}{2}}$

Q4. For an electric field $\vec{E} = k\sqrt{x}\hat{x}$ where k is a non-zero constant, total charge enclosed by the cube as shown below is



(a) 0

(b) $k \epsilon_0 l^{5/2} (\sqrt{3} - 1)$

(c) $k \epsilon_0 l^{5/2} (\sqrt{5} - 1)$

(d) $k \epsilon_0 l^{5/2} (\sqrt{2} - 1)$

Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

Q5. Consider a point particle A of mass m_A colliding elastically with another point particle

B of mass m_B at rest, where $\frac{m_B}{m_A} = \gamma$. After collision, the ratio of the kinetic energy of

particle B to the initial kinetic energy of particle A is given by

(a) $\frac{4}{\gamma + 2 + \frac{1}{\gamma}}$

(b) $\frac{2}{\gamma + \frac{1}{\gamma}}$

(c) $\frac{2}{\gamma + 2 - \frac{1}{\gamma}}$

(d) $\frac{1}{\gamma}$

Q6. Two classical particles are distributed among $N (> 2)$ sites on a ring. Each site can accommodate only one particle. If two particles occupy two nearest neighbour sites, then the energy of the system is increased by ϵ . The average energy of the system at temperature T is


(a) $\frac{2\epsilon e^{-\beta\epsilon}}{(N-3) + 2e^{-\beta\epsilon}}$

(b) $\frac{2N\epsilon e^{-\beta\epsilon}}{(N-3) + 2e^{-\beta\epsilon}}$

(c) $\frac{\epsilon}{N}$

(d) $\frac{2\epsilon e^{-\beta\epsilon}}{(N-2) + 2e^{-\beta\epsilon}}$

Q7. Consider a 741 operational amplifier circuit as shown below, where $V_{CC} = V_{EE} = +15V$ and $R = 2.2 k\Omega$. If $v_i = 2mV$, what is the value of v_o with respect to the ground?



(a) $-1mV$

(b) $-2mV$

(c) $-3mV$

(d) $-4mV$


Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

- Q8. The Fourier transform of the function $\frac{1}{x^4 + 3x^2 + 2}$ up to proportionality constant is
- (a) $\sqrt{2} \exp(-k^2) - \exp(-2k^2)$ (b) $\sqrt{2} \exp(-|k|) - \exp(-\sqrt{2}|k|)$
- (c) $\sqrt{2} \exp(-\sqrt{|k|}) - \exp(-\sqrt{2|k|})$ (d) $\sqrt{2} \exp(-\sqrt{2}k^2) - \exp(-2k^2)$
- Q9. If $\rho = \frac{\left[I + \frac{1}{\sqrt{3}} (\sigma_x + \sigma_y + \sigma_z) \right]}{2}$, where σ 's are the Pauli matrices and I is the identity matrix, then the trace of σ^{2017} is
- (a) 2^{2017} (b) 2^{-2017} (c) 1 (d) $\frac{1}{2}$
- Q10. A cylinder at temperature $T = 0$ is separated into two compartments A and B by a free sliding piston. Compartments A and B are filled by Fermi gases made of spin $1/2$ and $3/2$ particles respectively. If particles in both the compartments have same mass, the ratio of equilibrium density of the gas in compartment A to that of gas in compartment B is
- (a) 1 (b) $\frac{1}{3^{2/5}}$ (c) $\frac{1}{2^{2/5}}$ (d) $\frac{1}{2^{2/3}}$
- Q11. What is the DC base current (approximated to nearest integer value in μA) for the following $n - p - n$ silicon transistor circuit, given $R_1 = 75 \Omega, R_2 = 4.0 k\Omega, R_3 = 2.1 k\Omega, R_4 = 2.6 k\Omega, R_5 = 6.0 k\Omega, R_6 = 6.8 k\Omega, C_1 = 1 \mu F, C_2 = 2 \mu F, V_C = +15V, \beta_{dc} = 75$?



- (a) 20 (b) 24 (c) 16 (d) 32

Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

Q12. Consider a particle confined by a potential $V(x) = k|x|$, where k is a positive constant.

The spectrum E_n of the system, within the WKB approximation is proportional to

(a) $\left(n + \frac{1}{2}\right)^{3/2}$ (b) $\left(n + \frac{1}{2}\right)^{2/3}$ (c) $\left(n + \frac{1}{2}\right)^{1/2}$ (d) $\left(n + \frac{1}{2}\right)^{4/3}$

Q13. Consider the Hamiltonian

$$H(t) = \alpha \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix} + \beta t \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & -2 \end{pmatrix}$$

The time dependent function $\beta(t) = \alpha$ for $t \leq 0$ and zero for $t > 0$. Find

$\left| \langle \Psi(t < 0) | \Psi(t > 0) \rangle \right|^2$, where $|\Psi(t < 0)\rangle$ is the normalised ground state of the system at a time $t < 0$ and $|\Psi(t > 0)\rangle$ is the state of the system at $t > 0$.

(a) $\frac{1}{2}(1 + \cos(2\alpha t))$ (b) $\frac{1}{2}(1 + \cos(\alpha t))$
 (c) $\frac{1}{2}(1 + \sin(2\alpha t))$ (d) $\frac{1}{2}(1 + \sin(\alpha t))$

Q14. The function $f(x) = \cosh x$ which exists in the range $-\pi \leq x \leq \pi$ is periodically repeated between $x = (2m-1)\pi$ and $(2m+1)\pi$, where $m = -\infty$ to ∞ . Using Fourier series, indicate the correct relation at $x = 0$

(a) $\sum_{n=-\infty}^{\infty} \frac{(-1)^n}{1-n^2} = \frac{1}{2} \left(\frac{\pi}{\cosh \pi} - 1 \right)$ (b) $\sum_{n=-\infty}^{\infty} \frac{(-1)^n}{1-n^2} = 2 \frac{\pi}{\cosh \pi}$
 (c) $\sum_{n=-\infty}^{\infty} \frac{(-1)^{-n}}{1+n^2} = 2 \frac{\pi}{\sinh \pi}$ (d) $\sum_{n=1}^{\infty} \frac{(-1)^n}{1+n^2} = \frac{1}{2} \left(\frac{\pi}{\sinh \pi} - 1 \right)$

Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

Q15. A toy car is made from a rectangular block of mass M and four disk wheels of mass m and radii r . The car is attached to a vertical wall by a massless horizontal spring with spring constant k and constrained to move perpendicular to the wall. The coefficient of static friction between the wheel of the car and the floor is μ . The maximum amplitude of oscillations of the car above which the wheels start slipping is

(a)
$$\frac{\mu g (M + 2m)(M + 4m)}{mk}$$

(b)
$$\frac{\mu g (M^2 - m^2)}{Mk}$$

(c)
$$\frac{\mu g (M + m)^2}{2mk}$$

(d)
$$\frac{\mu g (M + 4m)(M + 6m)}{2mk}$$

Head office

fiziks, H.No. 40 D, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16